



Acer platanoides 'Reitenbachii'



Height	15-20 m
Width	10-12m
Crown	broad ovoid, dark, dense crown
Bark and branches	grey
Leaf	green, redbrown
Flowers	yellowgreen, flowers in April
Fruits	brown, inconspicuous
Spines/thorns	None
Toxicity	has toxic components
Soil type	clayed soil, loamy soil, sandy soil
Soil moisture	withstand short flood
Paving	tolerates partial paving
Winter hardiness zone	4 (-34,4 to -28,9 °C)
Wind resistance	not very sensitive, not resistant to sea wind
Other resistances	resistant to frost (WH 1 - 6)
Fauna tree	resistant to frost (WH 1 - 6), valuable for butterflies
Application	avenues and broad streets
Shape	clearstem tree

This tree's history goes back to the nursery of J. Reitenbach in Poland, at that time part of Russia. The plant was named by Johann Xaver Robert Caspary, a German botanist, but it was introduced to the market by the Belgian nursery of Louis Van Houtte. In terms of habit, 'Reitenbachii' is typical of the species. The sprouting leaf is a shiny dark brown, turning to matt dark green with dark red veins in summer. An attractive tree, somewhat resembling 'Schwedleri', which has more intense leaf colours. Grows on any soil, but does not tolerate stagnating groundwater and peaty soils.