



Quercus cerris 'Summer Veil'



Height	5 - 6 m
Width	3-4m
Crown	broad pyramidal, semi-open crown
Bark and branches	twigs grey-brown to grey-green, felt-like, bark almost black, roughly grooved
Leaf	Elongated to oval, lobes very deeply incised, green, 6 - 12cm
Flowers	long ovoid, 2.5 - 3 cm, 1 - 4 together, cupule with long, narrow recurved scales
Fruits	Tree nursery M.M. Bömer, Zundert, Netherlands, 1995
Spines/thorns	None
Toxicity	usually not toxic to people, (large) pets and livestock
Soil type	preferably calcareous
Soil moisture	suitable for wet soil
Paving	tolerates no paving
Winter hardiness zone	6a (-23,3 to -20,6 °C)
Wind resistance	good
Other resistances	resistant to frost (WH 1 - 6), can withstand wind, resistant to de-icing salt
Application	coastal areas, industrial areas
Shape	clearstem tree
Origin	Middelgrote boom met een ietwat onregelmatige breed kegelvormige tot ronde kroon. Groeit beduidend langzamer dan de soort en blijft uiteindelijk kleiner. De stam heeft minder diktegroei en de twijgen zijn relatief dun. Het meest opvallende kenmerk van 'Summer Veil' zijn de zeer diep gelobde bladeren. De bladeren zijn op veel plaatsen tot op de hoofdnerf gelobt. De puntige lobben zijn op hun beurt ook grof getand. De bladern zijn 's zomers groen, kleuren geelbruin in de herfst en blijven tot diep in de winter verdroogd in de boom hangen. Gevoelig voor bastscheuren bij vorst. De diep gelobde bladeren geven de boom een fraaie, fijne textuur. Deze cultivar ontstond als zaailing van Q. cerris en werd door Van den Berk Boomkwekerijen, Sint-Oedenrode, Nederland op de markt gebracht.

Medium-sized tree with a somewhat irregular broadly conical to round crown. Grows significantly slower than the species and eventually stays smaller. The trunk has less thickness growth and the twigs are relatively thin. The most striking feature of 'Summer Veil' is its very deeply lobed leaves. The leaves are lobed to the main vein in many places. In turn, the pointed lobes are also coarsely toothed. The foliage is green in summer, turning yellow-brown in autumn and remaining on the tree until deep into winter, when it dries out. Prone to bark cracking in frost. The deeply lobed leaves give the tree an attractive, fine texture. This cultivar originated as a seedling of Q. cerris and was marketed by Van den

